

The Impact of Climate Change on Regional Air Quality

Authors: Chris Nolte¹, Ellen Cooter¹, Robert Gilliam¹, William Benjey¹, Jenise Swall¹, Alice Gilliland¹, Loretta Mickley², Peter Adams³

¹Atmospheric Sciences Modeling Division/Air Resources Laboratory/National Oceanic and Atmospheric Administration, in partnership with the National Exposure Research Laboratory/Office of Research and Development/US EPA

²Department of Earth and Planetary Sciences and Division of Engineering and Applied Sciences, Harvard University

³Department of Civil and Environmental Engineering and Department of Engineering and Public Policy, Carnegie Mellon University

Keywords: air quality modeling, CMAQ, climate change, particulate matter, global climate model

The Climate Impact on Regional Air Quality (CIRAQ) project is an ongoing collaborative research effort involving multiple Federal agencies and academic institutions that assesses the impact of future climate change on US regional air quality. In this study, air quality is simulated using the Community Multiscale Air Quality (CMAQ) model under current climatological conditions and under climatological conditions 50 years in the future. A regional climate model (RCM) is driven by a global circulation model (GCM) under one proposed scenario of future greenhouse gas emissions, and a series of regional air quality model simulations is then performed to test the sensitivity of ozone and PM_{2.5} (fine particulate matter) concentrations to changes in regional climate. A preliminary evaluation of the model's ability to represent current climate and air quality is presented and compared with model predictions for the 2050 time period. For the current period, the model adequately simulates means and variability of key climatic variables (surface temperature, pressure, and wind speeds) over the western US, but fails to capture some important synoptic-scale features observed in the eastern US. Although predicted weather patterns remain largely unchanged from their current state under this scenario, modeled average temperatures for the future period are increased by 2–3 K in the Southwest and about 1 K elsewhere. Preliminary air quality modeling results suggest that, for the domain as a whole, the impact of these climatic changes on ozone and PM_{2.5} concentrations is modest, especially when considering nationwide seasonally- or annually-averaged timescales. Future analysis will examine regional and seasonal variations in air quality, which can be expected to be most pronounced in areas such as the Southwest US, where the model results indicate the greatest future warming.

Disclaimer: *The research presented here was performed under the Memorandum of Understanding between the U.S. Environmental Protection Agency (EPA) and the U.S. Department of Commerce's National Oceanic and Atmospheric Administration (NOAA) and under agreement number DW13921548. This work constitutes a contribution to the NOAA Air Quality Program. Although it has been reviewed by EPA and NOAA and approved for publication, it does not necessarily reflect their policies or views.*

Point of Contact:

Chris Nolte

Physical Scientist

U.S. EPA/Atmospheric Modeling Division (AMD)

E243-01

109 TW Alexander Dr.

Research Triangle Park, NC 27711

919-541-2652

nolte.chris@epa.gov